

## Quality Evaluation of Strawberry Juice with TSS of 30.5° Brix Preserved with Benzoate, Sorbate, and Antioxidant Stored At Refrigeration Temperature

Shakoor Wisal <sup>a</sup>

Alia Gul <sup>b</sup>

Hina Gul <sup>c</sup>

**Abstract:** *The aim of this research was carried out to investigate the effect of different sucrose concentrations and chemical preservatives and storage temperature, i.e., refrigeration temperature (4-10°C), on the physicochemical and sensory attributes of strawberry juice stored for one year. samples were numbered as,  $T_{R32}$ =Strawberry juice (30.5° brix) - no preservative,  $T_{R33}$ =Strawberry juice (30.5° brix) with 0.1% sodium benzoate,  $T_{R34}$ =Strawberry juice (30.5° brix) with 0.1% potassium sorbate,  $T_{R35}$ =Strawberry juice (30.5° brix) with 0.05% sodium benzoate + 0.05% potassium sorbate,  $T_{R44}$ =Strawberry juice (30.5° brix) - no preservatives (control),  $T_{R45}$ =Strawberry juice (30.5° brix) with 0.1% sodium benzoate,  $T_{R46}$ =Strawberry juice (30.5° brix) with 0.1% potassium sorbate,  $T_{R47}$ =Strawberry juice (30.5° brix) with 0.05% sodium benzoate + 0.05% potassium sorbate. As a result, a decrease in color was recorded for all samples, but it was in the acceptable range during storage. Flavor deterioration occurred in control samples ( $T_{R32}$  and  $T_{R44}$ ), while treated samples showed less loss in flavor during storage. Comparatively better consistency was recorded by treated juice. After one year of storage, the samples were acceptable to the consumer on the basis of color, flavor, consistency, and overall acceptability. Among all the treatments,  $T_{R32}$  and  $T_{R44}$  were rejected during storage, while  $T_{R47}$  was found most effective, followed by  $T_{R45}$ , and showed better storage stability than other samples.*

**Key Words:** Strawberry Juice, Benzoate, Sorbate, Sucrose, Refrigeration Temperature

### Introduction

Strawberry (*Fragaria* sp.) is a member of the herbaceous perennial of the "shifts" the family; it is the most important fruit among the berries and has and possesses an important place among small fruit plants (Sharma *et al.* 2009). It is thought that berries are important against several types of cancers (Navindra and P.Seeram, 2008). It is grown in many countries of the world but is grown widely in the United States, Japan, Mexico, Italy, and Lebanon (Sharma *et al.* 2009). Strawberries are also grown wildly with a pleasant flavor but are smaller in size (Ensminger *et al.* 2007). Gradually

grown in Pakistan because its scope in the future is bright for farmers, and fruits can be traced to the maximum economic benefit to farmers. There are certain factors that affect the production of strawberries; for example, scientifically grown strawberries give higher production than conventional cultivation (Tonture *et al.* 2009). Strawberries and cooling requirements are low and can be successfully grown crop in the tropics and subtropics. Mature fruit in a short period of time, which is 30 to 40 days, non-perishable and must be consumed early after winning. It cannot reach far away markets in fresh form, so it should be treated carefully during transportation.

<sup>a</sup> Department of Agriculture, Bacha Khan University Charsadda, KP, Pakistan. Email: [swisalaup.edu.pk](mailto:swisalaup.edu.pk) (Corresponding Author)

<sup>b</sup> Department of Botany, Hazara University Mansehra, KP, Pakistan.

<sup>c</sup> Department of Botany, Hazara University Mansehra, KP, Pakistan.

**Citation:** Wisal, S., Gul, A., & Gul, H. (2022). Quality Evaluation of Strawberry Juice with TSS of 30.5° Brix Preserved with Benzoate, Sorbate, and Antioxidant Stored At Refrigeration Temperature. *Global Drug Design & Development Review*, VII(1), 1-10. [https://doi.org/10.31703/gdddr.2022\(VII-I\).01](https://doi.org/10.31703/gdddr.2022(VII-I).01)

Strawberries are the first fresh fruit on the market during the spring. Because of the delicious flavor and attractive color and structure of the format, and consumer demand for fruit growing, not only in Pakistan but also in other parts of the world. And consumed mainly fruits and fresh, and are also used in the form of treatment such as canned, cooked, and sweetened, and that is, jam, jelly, and frozen whole berries. Several products are prepared from it, such as purees, jam, juice, wine, etc. (Sharma *et al.* 2009). The fruit is firm, red meat, and sweet. More than 50% of sucrose in strawberries is glucose. Fruit contains citric acid and often some malic acid. The red color of the fruit is due to the presence of anthocyanin pigment. Volatile esters responsible for many of the fruit flavor are found in it. Nutritionally, strawberries contain carbohydrates low-calorie and a potential source of vitamin C, fiber and provide more vitamin C than oranges. The main components of fruit, vitamin C in strawberries is (64.0 mg), water (91.75 g) and protein (0.61 g), fat (0.37 g) and carbohydrate (7.02 g), fiber (2.3 grams), calcium (14.0 mg) and potassium (166.0 mg/160 g), respectively, and vitamin (a) (27). International units, ranging in pH from 3.27 -3.86, help in achieving stability in the pH range of color from 0.58 to 1.35%, and citric acid and malic and organic acids contribute to excellent flavor. TSS is in the range of 8.0 up to 11.5% and is ideal for juice called for in the market. Soluble solids / acid ratios from 8.52 to 13.79 are a good balance of sweet-tart flavor. In the past decade, there has been a significant increase in the demand for strawberries because the juice has excellent flavor with an attractive color. However, being the result of a soft tight little fluctuation in temperature results in damage and waste of fruit. Due to the lack of other storage facilities to keep the product in the form of raw materials, this study has begun to take advantage of the fruit is very perishable in the form of juice and study the effect of chemical preservatives on the quality of conservation of strawberry juice stored at ambient temperature. Potassium sorbate is important in the preservation of juices during storage (Ayub *et al.* 2010). Sodium benzoate can maintain the good quality of fruit

juices during preservation (Ayub *et al.* 2010). It is hoped that these results will help beverage industries to take advantage of this fruit juice to prepare with the shelf stability and the growing demand for consumer goods. This offer high-yield income for farmers and will improve the economy of Pakistan.

## Materials and Methods

Fresh mature and sound strawberries were purchased from the local fruit market of Peshawar and were brought to the Food Processing and Analytical Laboratory of the Department of Food Science and Technology, The University of Agriculture Peshawar, Pakistan, where research work was carried out. The strawberries were washed, followed by sorting, the juice was extracted using a juice extracting machine. The samples were numbered as, TR<sub>32</sub>=Strawberry juice (30.5°brix) - no preservatives (control), TR<sub>33</sub>=Strawberry juice (30.5°brix) with 0.1% sodium benzoate, TR<sub>34</sub>=Strawberry juice (30.5°brix) with 0.1% potassium sorbate, TR<sub>35</sub>=Strawberry juice (30.5°brix) with 0.05% sodium benzoate + 0.05% potassium sorbate, TR<sub>44</sub>=Strawberry juice (30.5°brix) - no preservatives (control), TR<sub>45</sub>=Strawberry juice (30.5°brix) with 0.1% sodium benzoate, TR<sub>46</sub>=Strawberry juice (30.5°brix) with 0.1% potassium sorbate, TR<sub>47</sub>=Strawberry juice (30.5°brix) with 0.05% sodium benzoate + 0.05% potassium sorbate. The juice was filled in a glass bottle, sealed, and stored at refrigeration temperature (4 - 10°C) for a period of one year.

## Chemical Analysis

The total soluble solids (TSS) were determined by using an Abbe refractometer at ambient temperature (AOAC, 2000). Inolab digital pH meter was used for pH determination. Acidity was determined by dissolving a known weight of sample in distilled water and titration against 0.01 N NaOH using phenolphthalein as an indicator (Srivastava and Sanjeev, 2003). Ascorbic acid was determined by the direct colorimetric method using 2, 6- dichlorophenol- indophenols as a

decolorizing agent by ascorbic acid in sample extract and in slandered ascorbic acid solution (AOAC, 2000). Reducing and non-reducing sucrose was determined by the lane Eynon method (AOAC, 2000).

## Sensory Evaluation

A panel of ten judges selected from staff and students of the food science department evaluated the product fortnightly for color, flavor, consistency, and overall acceptability by the method of [Larmond \(1977\)](#) using a scale from 1 to 9, where 1 represented extremely disliked and 9 represent extremely liked.

## Statistical Analysis

The data obtained were subjected to statistical analysis using RCBD (Randomized Complete Block Design), and the means were compared by using LSD (Least Significant Difference) test ([Steel and Torrie, 1980](#)). For all the analyses, the alpha error was set at 0.05%.

## Results and Discussion

### Chemical Analysis

Our results indicated that storage period and the temperature had a significant effect on total soluble solids (TSS) during storage. The mean TSS values of all samples decreased from 30.50 to 30.38 during storage. Minimum and a similar increase occurred in  $T_{R35}$ ,  $T_{R45}$ ,  $T_{R46}$ , and  $T_{R47}$  (6.55%) while the maximum increase was observed in  $T_{R33}$  and  $T_{R34}$  (8.19%) during storage at refrigeration temperature. Decrease was observed in control samples ( $T_{R32}$  and  $T_{R44}$ ) during storage. These results are in agreement with the results obtained by [Zeb et al. \(2009\)](#) during the preservation of grape juice stored at room temperature for one month preserved with sodium benzoate and potassium sorbate. Similar results were obtained by [Hussain et al. \(2011\)](#) during storage of apple and apricot blended juice storage at refrigeration temperature for three months. An increase in total soluble solids may be due to the breakdown of polysaccharides into monosaccharides and

oligosaccharides, while a decrease may be due to fermentation of sugars into ethyl alcohol, carbon dioxide and water.

The mean pH values of all samples decreased from 3.66 to 3.26 during storage. During storage maximum decrease in pH content was observed in  $T_{R32}$  (16.30%) followed by  $T_{R44}$  (15.93%) while a minimum decrease was observed in  $T_{R47}$  (7.96%) followed by  $T_{R45}$  (8.51%). Similar results were recorded by [Mehmood et al. \(2008\)](#) during the study of the effect of pasteurization and chemical preservatives on the quality and shelf stability of apple juice stored at ambient temperature for three months. During storage of apple and apricot blended juices storage, preserved with sodium benzoate at refrigeration temperature for three months by [Hussain et al. \(2011\)](#), decrease in pH was recorded. Decrease in pH may be due to conversion of pectin into pectenic acid, which increases acidity and decreases pH of the juice.

The mean titratable acidity values of all samples increased from 0.95 to 1.84 during storage. Storage and treatments have a significant effect on the titratable acidity of strawberry juice. Maximum increase was observed in  $T_{R32}$  (124.69%) followed by  $T_{R44}$  (100.00%) while the minimum increase was observed in  $T_{R47}$  (78.18%) followed by  $T_{R45}$  (80.00%). Similarly increase in titratable acidity was observed by [Zeb et al. \(2009\)](#) during preservation of grape juice with sodium benzoate and potassium sorbate, stored at room temperature for one month storage period. The results of [Ayub and Bilal \(2001\)](#) are in agreement with our results, who observed an increase in acidity of pomegranate syrup, preserved under different light conditions and different packaging materials at room temperature for storage period of four months. This increase might be due to acidic compounds formed by degradation or oxidation of reducing sugar and high temperature. An increase in titratable acidity may be due to the breakdown of pectin into pectenic acid or due to the formation of acid by the breakdown of polysaccharides or oxidation of reducing sugars.

Ascorbic acid is sensitive to heat, light and oxygen etc and is the most difficult vitamin to be

preserved during storage. As it is the least stable vitamin, it decreases in the product during storage. The mean ascorbic acid values of all samples decreased from 24.45 to 9.05 during storage. Maximum decrease was observed in  $T_{R32}$  (79.55%) followed by  $T_{R44}$  (70.70%) while a minimum decrease was observed in  $T_{R47}$  (54.29%) followed by  $T_{R45}$  (57.04%). The results are in agreement with the findings of Zeb *et al.* (2009) during the preservation of grape juice with sodium benzoate and potassium sorbate, stored at room temperature for one month storage period, who observed decrease in ascorbic acid content of grape juice. The results of Ayub and Bilal (2001) are in agreement with our results, who observed a decrease in ascorbic acid content of pomegranate syrup, preserved under different light conditions and different packaging materials at room temperature for storage period of four months. The losses may be due to oxygen present in the product and the headspace of the package.

Sugars are the most important constituent of fruit products and are an essential factor for the flavor of the food product and also act as a natural food preservative. The mean reducing sugars values of all samples increased from 23.58 to 27.24 during storage. Results showed that reducing sugars increased during storage and maximum increase was observed in  $T_{R34}$  (30.70%) followed by  $T_{R33}$  (30.44%) while minimum increase was observed in  $T_{R47}$  (27.14 %) followed by  $T_{R45}$  (28.15%) while control samples i. e.  $T_{R32}$  and  $T_{R44}$  showed decrease during storage. Mehmood *et al.* (2008) observed an increase in reducing sugar of apple juice preserved with chemical preservatives stored at ambient temperature for three months. Increase in reducing sugar of apple and apricot blended juice, preserved with sodium benzoate at refrigeration temperature for three months, and was observed by Hussain *et al.* (2011). This increase in reducing sugar might be due to the conversion of sucrose to reducing sugars (glucose and fructose) primarily due to acids and higher temperature.

The mean non reducing sugars values of all samples decreased from 5.04 to 1.34 during

storage. Maximum decrease in non reducing sugar was observed in  $T_{R32}$  (93.65%) followed by  $T_{R44}$  (92.46%) while a minimum decrease was observed in  $T_{R47}$  (65.87%) followed by  $T_{R45}$  (66.27%). Mehmood *et al.* (2008) observed a decrease in non reducing sugar of apple juice preserved with chemical preservatives stored at ambient temperature for three months. An increase in reducing sugar of apple and apricot blended juice, preserved with sodium benzoate at refrigeration temperature for three months, was observed by Hussain *et al.* (2011). This decrease in non reducing sugar might be due to the conversion of sucrose to glucose and fructose, primarily due to an increase in acidity and high storage temperature and storage period length.

### Sensory Analysis

The analysis of our data showed that storage period and treatments had a significant effect on sensory attributes (color, flavor, consistency and overall acceptability) of the strawberry juice. Loss in color was recorded in all treatments in which maximum loss was recorded in control samples while treated samples retained better color during storage at refrigeration temperature. Off flavor was produced in control samples which may be due to fermentation process in which ethyl alcohol, carbon dioxide and water were produced while treated samples retained better flavor which may be due to the combined effect of preservatives, refrigeration temperature and sugar concentration. Better consistency was recorded by treated samples, especially by those samples which were sweetened with sugar. Data showed that all the samples were acceptable even after one year of storage period. The results are in agreement with the findings of Zeb *et al.* (2009) during preservation of grape juice with sodium benzoate and potassium sorbate, stored at room temperature for one month storage period, who observed decrease in color of grape juice. The results of Ayub and Bilal (2001) are in agreement with our results, who observed a decrease in flavor of pomegranate syrup, preserved under different light conditions and different packaging materials

at room temperature for a storage period of four months. These results are in agreement with findings of Nilugen and Mahendran (2010), who observed a decrease in consistency of ready to serve beverages prepared from palmyrah fruit pulp and stored at room temperature for six months.

## Conclusion

From this research study it was concluded that sugar concentration, combination of preservatives and storage temperature play a positive role in

extending the shelf life of strawberry juice. It was concluded that  $T_{R32}$ =Strawberry juice (30.5° brix) - no preservatives (control) and  $T_{R44}$ =Strawberry juice (30.5° brix) - no preservatives (control), were rejected while  $T_{R47}$ =Strawberry juice (30.5° brix) with 0.05% sodium benzoate +0.05% potassium sorbate maintained better quality followed by  $T_{R45}$ =Strawberry juice (30.5° brix) with 0.1% sodium benzoate stored at refrigeration temperature for one year period.

**Table 1.** Effect of Chemical Preservatives on TSS of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days											Mean
	Initial	20	40	60	80	120	180	240	300	360	% Inc/dec	
TSS (Total Soluble Solids) in juice												
$T_{R32}$	30.5	30.5	29.5	28.5	27.0	25.5	24.0	23.5	22.5	21.5	29.50*	26.30 c
$T_{R33}$	30.5	30.5	30.5	30.5	30.5	30.5	31.0	31.0	32.0	33.0	8.19	31.00 a
$T_{R34}$	30.5	30.5	31.0	31.0	31.5	31.5	31.5	32.5	32.5	33.0	8.19	31.55 a
$T_{R35}$	30.5	30.5	30.5	30.5	30.5	31.5	31.5	32.0	32.0	32.5	6.55	31.20 a
$T_{R44}$	30.5	30.5	30.5	29.5	29.5	28.5	27.5	27.0	27.0	25.5	16.39*	28.60 b
$T_{R45}$	30.5	30.5	30.8	31.0	31.5	32.0	32.0	32.0	32.5	32.5	6.55	31.23 a
$T_{R46}$	30.5	31.0	31.5	31.5	31.5	32.0	32.0	32.0	32.5	32.5	6.55	31.70 a
$T_{R47}$	30.5	30.5	31.0	31.5	31.5	32.0	32.0	32.0	32.5	32.5	6.55	31.60 a
Mean	30.50 a	30.56 a	30.66 a	30.50 a	30.44 a	30.44 a	30.19 a	30.25 a	30.44 a	30.38 a		

Values followed by different Letters are Significantly ( $p \leq 0.05$ ) different from each other.

**Table 2.** Effect of Chemical Preservatives on pH of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days											Mean
	Initial	20	40	60	80	120	180	240	300	360	% Dec	
pH value												
$T_{R32}$	3.68	3.57	3.45	3.38	3.35	3.32	3.28	3.21	3.14	3.08	16.30	3.34 b
$T_{R33}$	3.68	3.67	3.51	3.49	3.42	3.41	3.36	3.35	3.33	3.31	10.05	3.45 a
$T_{R34}$	3.68	3.73	3.56	3.53	3.48	3.41	3.38	3.35	3.32	3.31	10.05	3.47 a
$T_{R35}$	3.68	3.63	3.54	3.53	3.48	3.42	3.39	3.37	3.35	3.33	9.51	3.47 a
$T_{R44}$	3.64	3.45	3.41	3.36	3.32	3.29	3.24	3.18	3.11	3.06	15.93	3.30 c
$T_{R45}$	3.64	3.54	3.49	3.47	3.46	3.44	3.42	3.38	3.35	3.33	8.51	3.45 a

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Dec	Mean
T <sub>R46</sub>	3.64	3.56	3.51	3.48	3.47	3.45	3.43	3.39	3.35	3.32	8.79	3.46 a
T <sub>R47</sub>	3.64	3.56	3.52	3.48	3.46	3.45	3.43	3.40	3.37	3.35	7.96	3.46 a
Mean	3.66 a	3.58 b	3.49 c	3.46 cd	3.43 de	3.39 ef	3.36 fg	3.32 gh	3.29 hi	3.26 i		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 3.** Effect of Chemical Preservatives on Acidity (%) of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Inc	Mean
Acidity % content												
T <sub>R32</sub>	0.81	1.12	1.20	1.32	1.44	1.52	1.62	1.68	1.76	1.82	124.69	1.42 c
T <sub>R33</sub>	0.81	0.96	1.06	1.15	1.26	1.36	1.42	1.48	1.54	1.60	97.53	1.26 d
T <sub>R34</sub>	0.81	0.96	1.06	1.16	1.26	1.37	1.43	1.48	1.54	1.60	97.53	1.26 d
T <sub>R35</sub>	0.81	0.96	1.04	1.14	1.24	1.36	1.42	1.46	1.52	1.58	95.06	1.25 d
T <sub>R44</sub>	1.10	1.32	1.44	1.58	1.70	1.80	1.92	2.00	2.12	2.20	100.00	1.71 a
T <sub>R45</sub>	1.10	1.24	1.38	1.50	1.60	1.68	1.76	1.82	1.90	1.98	80.00	1.59 b
T <sub>R46</sub>	1.10	1.24	1.38	1.50	1.60	1.70	1.78	1.84	1.92	2.00	81.81	1.60 b
T <sub>R47</sub>	1.10	1.22	1.36	1.48	1.58	1.66	1.74	1.80	1.88	1.96	78.18	1.57 b
Mean	0.95 j	1.12 i	1.24 h	1.35 g	1.46 f	1.55 e	1.63 d	1.69 c	1.77 b	1.84 a		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 4.** Effect of Chemical Preservatives on Ascorbic Acid (%) of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Dec	Mean
Ascorbic acid content												
T <sub>R32</sub>	22.50	19.64	15.60	13.05	11.75	10.50	9.80	8.60	6.90	4.60	79.5 5	12.29f
T <sub>R33</sub>	22.50	19.50	17.20	15.88	13.88	12.32	11.50	9.98	8.40	7.32	67.4 6	13.85de
T <sub>R34</sub>	22.50	19.82	17.00	15.62	13.54	12.00	11.22	9.82	8.22	7.00	68.8 8	13.67e

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Dec	Mean
T <sub>R35</sub>	22.50	20.32	18.60	16.50	14.12	13.00	12.02	10.54	9.32	7.98	64.53	14.49d
T <sub>R44</sub>	28.40	23.40	18.70	15.22	13.76	12.48	11.36	10.40	9.76	8.32	70.70	15.18c
T <sub>R45</sub>	28.40	25.04	21.32	19.54	17.56	16.12	15.98	14.88	13.54	12.20	57.04	18.46ab
T <sub>R46</sub>	28.40	25.40	21.00	19.32	17.42	16.00	15.76	14.82	13.40	12.00	57.74	18.35b
T <sub>R47</sub>	28.40	26.72	22.04	20.50	18.12	16.88	16.02	15.34	14.00	12.98	54.29	19.10a
Mean	24.45 a	22.48 b	18.93 c	16.95 d	15.02 e	13.66 f	12.96 f	11.80 g	10.40 h	9.050 i		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 5.** Effect of Chemical Preservatives on Reducing Sugar of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Inc/dec	Mean
	Reducing sugar											
T <sub>R32</sub>	23.58	23.26	22.80	22.04	20.87	19.71	18.55	18.16	17.39	16.62	29.52*	20.30b
T <sub>R33</sub>	23.58	25.38	26.18	27.88	28.78	29.48	30.08	30.38	30.66	30.76	30.44	28.32a
T <sub>R34</sub>	23.58	25.40	26.22	27.94	28.84	29.54	30.14	30.42	30.72	30.82	30.70	28.36a
T <sub>R35</sub>	23.58	25.32	26.10	27.78	28.66	29.34	29.92	30.20	30.46	30.56	29.60	28.19 a
T <sub>R44</sub>	23.58	23.34	23.22	22.80	22.62	21.88	21.36	20.98	20.26	18.71	20.65*	21.88b
T <sub>R45</sub>	23.58	25.18	26.38	27.38	28.26	29.08	29.58	29.88	30.10	30.22	28.15	27.96a
T <sub>R46</sub>	23.58	25.20	26.40	27.40	28.28	29.10	29.60	29.90	30.12	30.24	28.24	27.98a
T <sub>R47</sub>	23.58	25.14	26.32	27.30	28.16	28.96	29.40	29.58	29.88	29.98	27.14	27.83a
Mean	23.58 c	24.78 bc	25.45 abc	26.32 ab	26.81 a	27.14 a	27.33 a	27.44 a	27.45 a	27.24 a		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 6.** Effect of Chemical Preservatives on Non-reducing Sugar of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treat. Ment	Storage period in days										% Dec	Mean
	Initial	20	40	60	80	120	180	240	300	360		
	Non-reducing sugar											
T <sub>R32</sub>	5.04	3.92	3.30	2.84	2.40	2.02	1.64	1.16	0.78	0.32	93.65	2.34 b
T <sub>R33</sub>	5.04	4.32	3.74	3.32	2.94	2.60	2.28	2.04	1.82	1.64	67.46	2.97 a
T <sub>R34</sub>	5.04	4.30	3.72	3.30	2.92	2.58	2.26	2.02	1.80	1.62	67.86	2.95 a
T <sub>R35</sub>	5.04	4.34	3.76	3.34	2.96	2.62	2.30	2.06	1.84	1.66	67.06	2.99 a
T <sub>R44</sub>	5.04	4.00	3.36	2.90	2.48	2.08	1.70	1.24	0.84	0.38	92.46	2.40 b
T <sub>R45</sub>	5.04	4.38	3.80	3.38	3.00	2.66	2.34	2.10	1.88	1.70	66.27	3.02 a
T <sub>R46</sub>	5.04	4.36	3.78	3.36	2.98	2.64	2.32	2.08	1.86	1.68	66.67	3.01 a
T <sub>R47</sub>	5.04	4.38	3.82	3.40	3.02	2.68	2.36	2.12	1.90	1.72	65.87	3.04 a
Mean	5.04 a	4.25 b	3.66 c	3.23 d	2.83 e	2.48 f	2.15 g	1.85 h	1.59 i	1.34 j		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 7.** Effect of Chemical Preservatives on Color of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days										% Dec	Mean
	Initial	20	40	60	80	120	180	240	300	360		
	Color score rate											
T <sub>R32</sub>	8.00	7.00	6.67	6.67	6.30	6.00	5.67	5.67	5.50	5.50	31.25	6.29 b
T <sub>R33</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 a
T <sub>R34</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 a
T <sub>R35</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 a
T <sub>R44</sub>	8.00	7.00	6.67	6.67	6.30	6.00	5.67	5.67	5.50	5.50	31.25	6.29 b
T <sub>R45</sub>	8.00	7.50	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.50	18.75	7.10 a
T <sub>R46</sub>	8.00	7.50	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.50	18.75	7.10 a
T <sub>R47</sub>	8.00	7.50	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.50	18.75	7.10 a
Mean	8.00 a	7.37 b	7.29 b	7.10 c	6.82 d	6.75 d	6.48 e	6.29 f	6.06 g	6.06 g		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 8.** Effect of Chemical Preservatives on Flavor of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days										% Dec	Mean
	Initial	20	40	60	80	120	180	240	300	360		
	Flavor score rate											
T <sub>R24</sub>	8.00	6.00	5.50	5.00	4.50	4.30	3.80	3.50	3.00	3.00	62.50	4.66 d
T <sub>R25</sub>	8.00	7.50	7.50	7.00	7.00	6.50	6.50	6.00	6.00	5.67	29.12	6.76 b
T <sub>R26</sub>	8.00	7.50	7.50	7.00	7.00	6.50	6.50	6.00	6.00	5.67	29.12	6.76 b
T <sub>R27</sub>	8.00	7.50	7.50	7.00	7.00	6.50	6.50	6.00	6.00	5.67	29.12	6.76 b



Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Dec	Mean
T <sub>R36</sub>	9.00	6.67	6.30	5.67	5.30	5.00	4.50	4.50	4.00	4.00	55.55	5.49 c
T <sub>R37</sub>	9.00	8.50	8.50	8.00	8.00	7.50	7.00	7.00	7.00	7.00	22.22	7.75 a
T <sub>R38</sub>	9.00	8.50	8.50	8.00	8.00	7.50	7.00	7.00	7.00	7.00	22.22	7.75 a
T <sub>R39</sub>	9.00	8.50	8.50	8.00	8.00	7.50	7.00	7.00	7.00	7.00	22.22	7.75 a
Mean	8.50	7.58	7.47	6.95	6.85	6.41	6.10	5.87	5.57	5.62		
	a	b	b	c	c	d	de	ef	ef	f		

Values followed by different Letters are Significantly ( $p \leq 0.05$ ) different from each other.

**Table 9.** Effect of Chemical Preservatives on Consistency of Strawberry Juice Stored at Refrigeration Temperature (4-10°C)

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Dec	Mean
	consistency score rate											
T <sub>R32</sub>	8.00	7.67	7.60	7.30	6.50	6.30	6.00	5.60	5.50	5.00	37.50	6.54 c
T <sub>R33</sub>	8.00	7.50	7.50	7.00	7.00	6.50	6.50	6.00	6.00	5.50	31.25	6.75 b
T <sub>R34</sub>	8.00	7.50	7.50	7.00	7.00	6.50	6.50	6.00	6.00	5.50	31.25	6.75 b
T <sub>R35</sub>	8.00	7.50	7.50	7.00	7.00	6.50	6.50	6.00	6.00	5.50	31.25	6.75 b
T <sub>R44</sub>	8.00	7.67	7.60	7.30	6.50	6.30	6.00	5.60	5.50	5.00	37.50	6.54 c
T <sub>R45</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.25	6.25	21.87	6.95 a
T <sub>R46</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 ab
T <sub>R47</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.25	6.25	21.87	6.95 a
Mean	8.00 a	7.54	7.52	7.07	6.87	6.63	6.37	6.08	5.93	5.62		
		b	b	c	c	d	e	f	f	g		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

**Table 10.** Effect of Chemical Preservatives on overall Acceptability of Strawberry Juice Stored at Refrigeration Temperature (4-10°C).

Treatment	Storage period in days											
	Initial	20	40	60	80	120	180	240	300	360	% Dec	Mean
	Overall acceptability score rate											
T <sub>R32</sub>	8.00	7.00	7.67	6.30	6.00	5.67	5.30	5.00	4.50	4.00	50.00	5.84 b
T <sub>R33</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	5.67	29.12	6.86 a
T <sub>R34</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	5.67	29.12	6.86 a
T <sub>R35</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	5.67	29.12	6.86 a
T <sub>R44</sub>	8.00	7.00	7.67	6.30	6.00	5.67	5.30	5.00	4.50	4.50	43.75	5.89 b
T <sub>R45</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 a
T <sub>R46</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 a
T <sub>R47</sub>	8.00	7.50	7.50	7.00	7.00	7.00	6.50	6.50	6.00	6.00	25.00	6.90 a
Mean	8.00 a	7.37	7.29	6.82	6.75	6.66	6.20	6.12	6.62	5.43		
		b	b	c	c	c	d	d	e	e		

Values followed by different letters are significantly ( $p \leq 0.05$ ) different from each other.

\* = decrease occurred in it.

## References

- Amin, N. U. (1996). Evaluation of different strawberry cultivars for offseason production under plastic tunnel.
- Ayub, M. J., Ullah, A., Muhammad, & Zeb, A. (2010). Evaluation of strawberry juice preserved with chemical preservatives at refrigeration temperature. *Int J. of Nutrition and Metab*, 2(2), 27-32.
- Ayub, M., & Khan, M. B. (2001). Effect of different light conditions and colored glass bottles on the retention of quality characteristics of pomegranate syrup during storage at room temperature. *Sarhad J. of Agric*, 17(4), 629-632.
- Hussain, I., Zeb, A., & Ayub. (2011). Evaluation of Apple and Apricot Blend Juice Preserved with Sodium Benzoate at Refrigeration Temperature. *World J. of Agric. Sci.*, 7(2), 136-142.
- Larmond, E. (1977). Method for sensory evaluation of food. *Canada Dept. of Agriculture, Publ.*, 1286, 36-37.
- Mehmood, Z., Zeb, A., Ayub, M., Bibi, N., Badshah, A., & Ihsanullah. (2008). Effect of Pasteurization and Chemical preservatives on the Quality and shelf stability of Apple Juice. *Amer. J. Food Tech*, 3(2), 147-153.
- Navindra, P., & Seeram. (2008). Berry Fruits for Cancer Prevention: Current Status and Future Prospects. *J. Agric. Food Chem*, 56, 630-635.
- Nilugin, S. E., & Mahendran, T. (2010). Preparation of ready-to-serve (RTS) beverage from palmyra (*Borassus flabellifer* L.) fruit pulp. *Journal of Agricultural Sciences*, 5(2), 234-237.
- Sharma, S., Joshi, V. K., & Abrol, G. (2009). An overview on strawberry [*Fragaria x ananassa* (Weston) Duchesne ex Rozier] wine production technology, composition, maturation and quality evaluation. *Natural product Radiance*, 8(4), 356-365.
- Srivastava, R. P., Sanjeev, K. (2003). Fruit and vegetable preservation principles and practices: Important methods for analysis of Fruits? Vegetables and their products. Third Revised and Enlarge Edition. International Book Distribution Co., Lucknow, India, 363.
- Steel, R. G. D., & Torrie, J. H. (1980). *Principle and procedures of statistics*. Mc Graw Hill Book Co. NY, U.S.A, 2nd ed, 195-238.
- Tönutare, T., Moor, U., Mölder, K., & Pöldma, P. (2009). Fruit composition of organically and conventionally cultivated strawberry "Polka". *Agronomy Research (Special Issue II)*, 755-7660.
- U.S Dept. of Agriculture USDA. (1998). Nutrient database for standard reference, (www Oregon-strawberries. Org).
- Zeb, A. I., Ahmad, K., & Ayub, M. (2009). Grape juice preservation with benzoate and sorbate. *J. Advance in Food Sci*, 31(1), 17-21.